

**§ 147.23 Hatchery sanitation.**

An effective program for the prevention and control of *Salmonella* and other infections should include the following measures:

(a) An effective hatchery sanitation program should be designed and implemented.

(b) The hatchery building should be arranged so that separate rooms are provided for each of the four operations: Egg receiving, incubation and hatching, chick/poult processing, and egg tray and hatching basket washing. Traffic and airflow patterns in the hatchery should be from clean areas to dirty areas (i.e., from egg room to chick/poult processing rooms) and should avoid tracking from dirty areas back into clean areas.

(c) The hatchery rooms, and tables, racks, and other equipment in them should be thoroughly cleaned and disinfected frequently. All hatchery wastes and offal should be burned or otherwise properly disposed of, and the containers used to remove such materials should be cleaned and sanitized after each use.

(d) The hatching compartments of incubators, including the hatching trays, should be thoroughly cleaned and disinfected after each hatch.

(e) Only clean eggs should be used for hatching purposes.

(f) Only new or cleaned and disinfected egg cases should be used for transportation of hatching eggs. Soiled egg case fillers should be destroyed.

(g) Day-old chicks, poults, or other newly hatched poultry should be distributed in clean, new boxes and new chick papers. All crates and vehicles used for transporting birds should be cleaned and disinfected after each use.

[67 FR 8474, Feb. 25, 2002]

**§ 147.24 Cleaning and disinfecting.**

The following procedures are recommended:

(a) In the poultry houses:

(1) Remove all live “escaped” and dead birds from the building. Blow dust from equipment and other exposed surfaces. Empty the residual feed from the feed system and feed pans and remove it from the building. Disassemble feeding equipment and dump and scrape as

needed to remove any and all feed cake and residue. Clean up spilled feed around the tank and clean out the tank. Rinse down and wash out the inside of the feed tank to decontaminate the surfaces and allow to dry.

(2) Remove all litter and droppings to an isolated area where there is no opportunity for dissemination of any infectious disease organisms that may be present. Housing where poultry infected with a mycoplasmal disease were kept should remain closed for 7 days before removal of the litter.

(3) Wash down the entire inside surfaces of the building and all the installed equipment such as curtains, ventilation ducts and openings, fans, fan housings and shutters, feeding equipment, watering equipment, etc. Use high pressure and high volume water spray (for example 200 pounds per square inch and 10 gallons per minute or more) to soak into and remove the dirt to decontaminate the building. Scrub the walls, floors, and equipment with a hot soapy water solution. Rinse to remove soap.

(4) Spray with a disinfectant which is registered by the Environmental Protection Agency as germicidal, fungicidal, pseudomonocidal, and tuberculocidal, in accordance with the specifications for use, as shown on the label of such disinfectant.

(b) In the hatchers and hatchery rooms:

(1) Use cleaning agents and sanitizers that are registered by the U.S. Environmental Protection Agency as germicidal, fungicidal, pseudomonocidal, and tuberculocidal. Use manufacturer's recommended dilution. Remove loose organic debris by sweeping, scraping, vacuuming, brushing, or scrubbing, or by hosing surface with high pressure water (for example 200 pounds per square inch and 10 gallons per minute or more). Remove trays and all controls and fans for separate cleaning. Use hot water (minimum water temperature of 140 °F) for cleaning hatching trays and chick separator equipment. Thoroughly wet the ceiling, walls, and floors with a stream of water, then scrub with a hard bristle brush. Use a cleaner/sanitizer that can penetrate protein and fatty deposits. Allow the chemical to cling to treated